GW150914: FACTSHEET

BACKGROUND IMAGES: TIME-FREQUENCY TRACE (TOP) AND TIME-SERIES (BOTTOM) IN THE TWO LIGO DETECTORS; SIMULATION OF BLACK HOLE HORIZONS (MIDDLE-TOP), BEST FIT WAVEFORM (MIDDLE-BOTTOM)

first direct detection of gravitational waves (GW) and first direct observation of a black hole binary

observed by	LIGO L1, H1	duration from 30 Hz	~ 200 ms
source type	black hole (BH) binary	# cycles from 30 Hz	~10
date	14 Sept 2015	peak GW strain	1 x 10 ⁻²¹
time	09:50:45 UTC	peak displacement of	±0.002 fm
likely distance	0.75 to 1.9 Gly 230 to 570 Mpc	interferometers arms frequency/wavelength	150 Hz, 2000 km
redshift	0.054 to 0.136	at peak GW strain peak speed of BHs	~ 0.6 c
signal-to-noise ratio	24	peak GW luminosity	3.6 x 10 ⁵⁶ erg s ⁻¹
false alarm prob.	< 1 in 5 million	radiated GW energy	2.5-3.5 M⊙
false alarm rate	< 1 in 200,000 yr	remnant ringdown fre	g. ~ 250 Hz
Source Masses M⊙		remnant damping tin	•
total mass	60 to 70	remnant size, area	180 km, 3.5 x 10 ⁵ km ²
primary BH	32 to 41	consistent with	passes all tests
secondary BH	25 to 33	general relativity?	performed
remnant BH	58 to 67	graviton mass bound	< 1.2 x 10 ⁻²² eV
mass ratio	0.6 to 1	coalescence rate of	
primary BH spin	< 0.7	binary black holes 2 to 400 Gp	2 to 400 Gpc ⁻³ yr ⁻¹
secondary BH spin	< 0.9	online trigger latency	~ 3 min
remnant BH spin	0.57 to 0.72	# offline analysis pipeli	
signal arrival time delay	arrived in L1 7 ms before H1	CPU hours consumed	~ 50 million (=20,000 PCs run for 100 days)
likely sky position	Southern Hemisphere	papers on Feb 11, 2016	13
likely orientation resolved to	face-on/off ~600 sq. deg.	# researchers	~1000, 80 institutions in 15 countries

Detector noise introduces errors in measurement. Parameter ranges correspond to 90% credible bounds. Acronyms: L1=LIGO Livingston, H1=LIGO Hanford; Gly=giga lightyear=9.46 x 10¹² km; Mpc=mega parsec=3.2 million lightyear, Gpc=10³ Mpc, fm=femtometer=10⁻¹⁵ m, M⊙=1 solar mass=2 x 10³⁰ kg